

Rapid Rat Immunoglobulin Isotyping Kit (Cat#: ISO-R8-IgG)

5 minute determination of 4 Rat Ig heavy chain isotypes

The Rapid Rat Immunoglobulin Isotyping Kit is a five-minute lateral flow assay with ELISA sensitivity for monoclonal antibody class and subclass determination. Acceptable sample types include hybridoma culture supernatant and purified monoclonal antibody. Ascites fluid is not recommended since it usually contains multiple immunoglobulins. However, this kit can still be used to determine the relative distribution of the main isotypes in a sample of ascites fluid.

Kit Components

The isotyping kit has two different sizes with 10 or 20 pouches containing two same cassettes per pouch, determining rat IgG1, IgG2a, IgG2b, and IgG2c. The strip contains a control line.

Sample Requirements and Dilution Conditions

Cell Culture Supernatant	Purified Antibody	Ascites
Recommended dilution is 1:10~1:100 in PBS.	Dilute samples to 0.5 - 2 µg/mL in PBS	Ascites fluid is not recommended since it usually contains multiple Igs. However, this kit can still be used to determine the relative distribution of the main isotypes in a sample of ascites fluid. The recommended dilution is 1:10,000 in PBS.

Protocol

- 1) Place the cassettes in horizontal plane.
- 2) Add 80-100 µL antibody solution directly to the Sample Well. Wait for 5 minutes.
- 3) A red test line at the location corresponding to antibody-specific isotype and a red control line at "C" position will appear within 1-5 minutes.

NOTE: Do not read after 20 minutes.

NOTE: If the color of tissue culture medium causes high background, 40-60 µL PBS can be chased to clear the background.

NOTE: Promptly return unused strips in their original bag and keep dry.

Result Interpretation

1. Invalid result: no red line at "C".
2. Negative result: only one red line at "C".
3. Positive results:
 - a). Single monoclonal antibody: In addition to the positive control "C" line, only one heavy chain and one light chain will appear.
 - b). Multiple monoclonal antibodies: In addition to the positive control line, more than one heavy chain and/or one light chain will appear. The line intensity can be used to judge the dominant isotypes.

Performance

Minimum detectable concentration is 100 ng/mL. There is no cross-reactivity among each isotype.

Storage

Store the strips in their original packaging at room temperature. This product is stable for up to 18 months from the certified date. Freezing the strips is not recommended.

Examples

